Double ratio of charmonia in p + Pb collisions at $\sqrt{s_{\rm NN}} = 5.02$ TeV

Yunpeng Liu[a], Che Ming Ko[a,b], Taesoo Song[a,c]

[a]*Cyclotron Institute, Texas A&M University, College Station, Texas 77843, USA*
[b]*Department of Physics and Astronomy, Texas A&M University, College Station, Texas 77843, USA*
[c]*Frankfurt Institute for Advanced Studies, J. W. Goethe University, 60438 Frankfurt am Main, Germany*

arXiv:1407.3805v1 [nucl-th] 14 Jul 2014

Abstract

Based on a kinetic description of J/ψ dissociation and production in an expanding quark-gluon plasma that is described by a 2+1 dimensional ideal hydrodynamics, we have studied the double ratio $R_{p\rm Pb}(\psi')/R_{p\rm Pb}(J/\psi)$ of charmonia in p + Pb collisions at $\sqrt{s_{\rm NN}} = 5.02$ TeV by including not only the cold nuclear matter effects but also the hot nuclear matter effects. We find that the double ratio of prompt charmonia is significantly suppressed in the most central collisions as a result of the hot nuclear matter effects.

Keywords: quark-gluon plasma, quarkonium suppression
PACS: 25.75.-q, 24.85.+p

Charmonia suppression in heavy ion collisions was first suggested in Ref. [1] based on the consideration of color screening in quark-gluon plasma (QGP). The anomalous suppression of J/ψ production, besides the normal suppression due to the cold nuclear matter effects, observed in Pb+Pb collisions at the Super Proton Synchrotron (SPS) seemed to confirm this prediction and thus suggested that a QGP was produced in these collisions [2]. Since then, there have been extensive experimental and theoretical studies on J/ψ production and suppression in both elementary and nuclear collisions. In particular,

- Baseline study of charmonia production in $p + p$ collisions has been carried out using the color singlet model (CSM) [3] and/or the color octet model (COM) [4].
- Cold nuclear matter effects, including the shadowing effect, the Cronin effect and the nuclear absorption, have been studied in p+A and d+A collisions.
- Hot medium effects, including the color screening, gluon dissociation, quasi-free scattering dissociation, regeneration and so forth, have been studied in $A + A$ collisions.

However, with the high beam energy at the LHC, the initial energy density in p+Pb collisions can be sufficient large for the formation of a QGP. It is then of interest to study the hot medium effects on quarkonia production in these collisions and compare them to those due to the cold nuclear matter [5–12].

In Ref. [13], we have carried out such a study by using a 2+1 dimensional ideal hydrodynamic model to describe the bulk dynamics of p + Pb collisions at $\sqrt{s_{\rm NN}} = 5.02$ TeV, i.e., based on the equation

$$\partial_\mu T^{\mu\nu} = 0 \quad (1)$$

with the boost invariant condition. In the above, $T^{\mu\nu} = (\epsilon + p)u^\mu u^\nu - g^{\mu\nu}p$ is the energy-momentum tensor in terms of the energy density ϵ, pressure p, and four velocity u. For the equation of state $\epsilon(p)$, we have used that of Ref. [14] based on an ideal gas of massless partons for the QGP and massive hadrons for the hadronic matter, together with a bag constant which leads to a critical temperature $T_c = 165$ MeV [15–17] at zero baryon chemical potential. For the initial energy density at time $\tau_0 = 0.6$ fm/c and its rapidity dependence, it is obtained from a multiple phase transport (AMPT) model [18]. Selecting the most central 10% collisions according to the yield of partons in AMPT, the maximum initial energy density is found at rapidity $y = -2$.

In Fig. 1, we show the energy density distribution in the transverse plane at rapidity $y = -2$ for different proper time τ. The origin of the coordinates is located at the center of the proton, and the center of the lead lies on the negative x-axis. The initial energy density is roughly isotropic in the transverse plane with a slightly larger value on the lead side. The maximum value is



Figure 1: (Color online) Energy density ϵ at rapidity $y = -2$ in the most central 10% p + Pb collisions at $\sqrt{s_{\rm NN}} = 5.02$ TeV as a function of the transverse coordinate x and y relative to the center of the proton at different proper time τ.

$\epsilon_0 \approx 15$ GeV/fm^3, which corresponds to a temperature $T_0 \approx 290$ MeV.

Since the energy density is large and the volume is small compared with those in A+A collisions at lower energies, the system expands fast, leading to a quick decreas of the energy density at the center. At time $\tau = 2.0$ fm/c, the energy density at the center is even smaller than that in the peripheral region, and the maximum energy density is at the radius $r \approx 1.5$ fm. The value of the energy density on the positive axis is slightly larger than that at the negative axis. Since the center of the QGP produced in p+Pb collisions hadronizes earlier, its spacial topology differs from that in Pb+Pb collisions[1].

For the time evolution of the distribution function $f(\mathbf{x}, \mathbf{p}, t)$ of J/ψ in the phase space of coordinate $\mathbf{x}$ and momentum $\mathbf{p}$, we use the transport equation

$$\partial_t f + \mathbf{v} \cdot \nabla f = -\alpha f + \beta, \tag{2}$$

where $\mathbf{v}$ is the velocity of the J/ψ, and α and β are the dissociation and the regeneration rate, respectively. The dissociation rate

$$\alpha(T, u, p) = \frac{N_g}{E} \int \frac{d\mathbf{k}}{(2\pi)^3 E_g} k \cdot p \, f_g(T, u, k) \sigma_D. \tag{3}$$

takes into account the gluon dissociation process $J/\psi + g \to c + \bar{c}$ with the cross section σ_D [14, 19, 20]. The regeneration rate takes into account the inverse process $c + \bar{c} \to J/\psi + g$, whose cross section is related to that of the gluon dissociation process by the detailed balance relation, and can be calculated using the distribution functions of charm and anticharm quarks. For simplicity, we take the charm quark distribution to be in thermal equilibrium, i.e.,

$$f_c(\mathbf{x}, \mathbf{q}_c) = \rho_c(\mathbf{x}) \frac{N}{e^{q_c \cdot u/T} + 1}, \tag{4}$$

where ρ_c is the number density of charm quarks and satisfies the conservation equation $\partial_t \rho_c + \nabla \cdot (\rho_c \mathbf{v}_m) = 0$ with $\mathbf{v}_m$ being the velocity of the medium, and $N \equiv \left[(2\pi)^{-3} \int d\mathbf{q}_c \left(e^{q_c \cdot u/T} + 1 \right)^{-1} \right]^{-1}$ is the normalization factor for the Fermi distribution. For anticharm quarks, their distribution function $f_{\bar{c}}$ has a similar form. We also include a velocity-dependent dissociation temperature T_d to describe the color screening effect [21].

Different from Pb+Pb collisions at the LHC, at most one pair of charm and anticharm quarks are produced in a p+Pb collision, and they are also not likely to reach thermal equilibrium . These effects are included via a canonical enhancement factor $C_{ce} = 1 + 1/(dN_c^{\rm dir}/dy)$ in the evaluation of the equilibrium number of J/ψ, as in the statistical model [22–24], and the relaxation reduction factor $r = 1 - \exp(-\tau/\tau_r)$ [25] with the charm quark relaxation time $\tau_r = 7$ fm/c [26, 27]. More details can be found in Ref. [13]. We note that these effects can be more accurately studied in a transport model [28].

For the initial distribution of J/ψ, it is obtained from the Glauber model with the inclusion of initial-state cold nuclear matter effects and a J/ψ production cross section of $d\sigma_{pp}/dy = 5.68\,\mu$b for pp collisions at 5.02 TeV. The latter is obtained from interpolating the experimental results at lower ($\sqrt{s_{\rm NN}} = 2.76$ TeV) [29] and higher ($\sqrt{s_{\rm NN}} = 7$ TeV) [30] energies with a power-law form and an average transverse momentum square $\langle p_T^2 \rangle = 9.5$ GeV^2 at middle rapidity. As to the rapidity distribution, it is assumed to be Gaussian with parameters taken from Refs. [31, 32]. The density distribution of initially produced J/ψ in space is assumed to be proportional to the thickness of a uniform solid sphere of radius $r = 0.8$ fm as that for the proton.

For the cold nuclear matter effects on J/ψ and charm quarks, they are assumed to be the same and are taken from Refs. [20, 33], which then lead to a suppression in proton rapidity and an enhancement in Pb rapidity.

The above treatment for the ground state J/ψ is generalized to include its excited states χ_c and ψ' with different dissociation temperatures T_D and cross sections [34, 35] as in Ref. [21]. For the initial abundance

[1]The Euler characteristic χ changes from 2 to 4 in central p+Pb collisions.

of these excited states, they are determined from that of J/ψ by using the empirically known feed-down contributions to J/ψ production in $p+p$ collisions, i.e., using the proportion 6 : 3 : 1 for direct J/ψ, feed-down from χ_c and from ψ' [36]. They are further assumed to subject to the same cold nuclear matter effects as the J/ψ.

For the contribution to J/ψ production from the decay of regenerated χ_c and ψ', they are included with the branch ratios from Ref. [37].



Figure 2: Double ratio $R^{\rm pro}_{p{\rm Pb}}(\psi')/R^{\rm pro}_{p{\rm Pb}}(J/\psi)$ of prompt charmonia in p + Pb collisions at $\sqrt{s_{\rm NN}}$ = 5.02 TeV as a function of rapidity for centralities 0-10% and 40-60%.

In Fig. 2, we show the double ratio of prompt charmonia as a function of rapidity. It is seen that the double ratio at rapidity from -4 to 0 is smaller than unit due to stronger suppression of ψ' in the hot medium. In the most central 10% collisions, it is smaller than 0.1, while in collisions with centrality 40%-60%, it is between 0.3 and 0.4. Since the formation time of J/ψ and ψ' is much longer than that for the proton and Pb to pass through each other, the cold nuclear matter effects are not expected to differ much between the J/ψ and ψ'.

In experiments, the inclusive double ratio is easier to measure. According to a recent study [38], the contribution of B meson decay increases the double ratio in heavy ion collisions. The corresponding result for p+Pb collisions is shown in Fig. 3. The qualitative effect of B meson decay is the same as that in Pb+Pb collisions [38], and the double ratio is above 0.4 in the whole rapidity range.

In summary, we have studied the double ratio of quarkonia in p + Pb collisions at $\sqrt{s_{\rm NN}}$ = 5.02 TeV in a transport approach based on a 2+1 dimensional ideal hydrodynamic model and found that both the prompt and inclusive double ratios are smaller than one, espe-



Figure 3: Double ratio $R^{\rm incl}_{p{\rm Pb}}(\psi')/R^{\rm incl}_{p{\rm Pb}}(J/\psi)$ of inclusive charmonia in p + Pb collisions at $\sqrt{s_{\rm NN}}$ = 5.02 TeV as a function of rapidity for centrality 40-60%.

cially for the prompt one in the most central 10% collisions. This is not expected from the cold nuclear matter effects as they exist before J/ψ and ψ' are formed. Measuring the double ratio of charmonia thus provides the possibility to study the hot medium effects in p+Pb collisions at the LHC.

Acknowledgements

This work was supported by the U.S. National Science Foundation under Grant No. PHY-1068572, the US Department of Energy under Contract No. DE-FG02-10ER41682, and the Welch Foundation under Grant No. A-1358.

References

[1] T. Matsui, H. Satz, J/psi Suppression by Quark-Gluon Plasma Formation, Phys. Lett. B178 (1986) 416. doi:10.1016/0370-2693(86)91404-8.

[2] M. Gonin, et al., Anomalous J / psi suppression in Pb + Pb collisions at 158-A-GeV/c, Nucl. Phys. A610 (1996) 404c–417c. doi:10.1016/S0375-9474(96)00373-9.

[3] C.-H. Chang, Hadronic Production of J/ψ Associated With a Gluon, Nucl. Phys. B172 (1980) 425–434. doi:10.1016/0550-3213(80)90175-3.

[4] G. T. Bodwin, E. Braaten, G. P. Lepage, Rigorous QCD analysis of inclusive annihilation and production of heavy quarkonium, Phys. Rev. D51 (1995) 1125–1171. arXiv:hep-ph/9407339, doi:10.1103/PhysRevD.55.5853,10.1103/PhysRevD.51.1125.

[5] C. Gerschel, J. Hufner, A Contribution to the Suppression of the J/psi Meson Produced in High-Energy Nucleus Nucleus Collisions, Phys. Lett. B207 (1988) 253–256. doi:10.1016/0370-2693(88)90570-9.

[6] T. Gousset, H. Pirner, The Ratio of gluon distributions in Sn and C, Phys. Lett. B375 (1996) 349–354. arXiv:hep-ph/9601242, doi:10.1016/0370-2693(96)00189-X.

[7] L. D. McLerran, R. Venugopalan, Fock space distributions, structure functions, higher twists and small x, Phys. Rev. D59 (1999) 094002. arXiv:hep-ph/9809427, doi:10.1103/PhysRevD.59.094002.

[8] B. Kopeliovich, B. Zakharov, Quantum effects and color transparency in charmonium photoproduction on nuclei, Phys. Rev. D44 (1991) 3466–3472. doi:10.1103/PhysRevD.44.3466.

[9] B. Kopeliovich, J. Nemchick, N. N. Nikolaev, B. Zakharov, Decisive test of color transparency in exclusive electroproduction of vector mesons, Phys. Lett. B324 (1994) 469–476. arXiv:hep-ph/9311237, doi:10.1016/0370-2693(94)90225-9.

[10] B. Kopeliovich, Transparent nuclei and deuteron gold collisions at RHIC, Phys. Rev. C68 (2003) 044906. arXiv:nucl-th/0306044, doi:10.1103/PhysRevC.68.044906.

[11] E. Iancu, R. Venugopalan, The Color glass condensate and high-energy scattering in QCDarXiv:hep-ph/0303204.

[12] E. Ferreiro, F. Fleuret, J. Lansberg, A. Rakotozafindrabe, Cold nuclear matter effects on J/psi production: Intrinsic and extrinsic transverse momentum effects, Phys. Lett. B680 (2009) 50–55. arXiv:0809.4684, doi:10.1016/j.physletb.2009.07.076.

[13] Y. Liu, C. M. Ko, T. Song, Hot medium effects on J/ψ production in $p + Pb$ collisions at $\sqrt{s_{NN}}$ = 5.02 TeV, Phys. Lett. B728 (2014) 437–442. arXiv:1309.5113, doi:10.1016/j.physletb.2013.12.016.

[14] Y.-p. Liu, Z. Qu, N. Xu, P.-f. Zhuang, J/psi Transverse Momentum Distribution in High Energy Nuclear Collisions at RHIC, Phys. Lett. B678 (2009) 72–76. arXiv:0901.2757, doi:10.1016/j.physletb.2009.06.006.

[15] Y. Aoki, Z. Fodor, S. Katz, K. Szabo, The QCD transition temperature: Results with physical masses in the continuum limit, Phys. Lett. B643 (2006) 46–54. arXiv:hep-lat/0609068, doi:10.1016/j.physletb.2006.10.021.

[16] S. Borsanyi, et al., Is there still any T_c mystery in lattice QCD? Results with physical masses in the continuum limit III, JHEP 1009 (2010) 073. arXiv:1005.3508, doi:10.1007/JHEP09(2010)073.

[17] Y. Aoki, G. Endrodi, Z. Fodor, S. Katz, K. Szabo, The Order of the quantum chromodynamics transition predicted by the standard model of particle physics, Nature 443 (2006) 675–678. arXiv:hep-lat/0611014, doi:10.1038/nature05120.

[18] Z.-W. Lin, C. M. Ko, B.-A. Li, B. Zhang, S. Pal, A Multi-phase transport model for relativistic heavy ion collisions, Phys. Rev. C72 (2005) 064901. arXiv:nucl-th/0411110, doi:10.1103/PhysRevC.72.064901.

[19] M. E. Peskin, Short Distance Analysis for Heavy Quark Systems. 1. Diagrammatics, Nucl. Phys. B156 (1979) 365. doi:10.1016/0550-3213(79)90199-8.

[20] L. Manceau, Quarkonium measurements in Pb-Pb and p-Pb collisions with ALICE at the LHCarXiv:1307.3098.

[21] Y. Liu, N. Xu, P. Zhuang, Velocity Dependence of Quarkonium Dissociation Temperature in High-Energy Nuclear Collisions, Phys. Lett. B724 (2013) 73–76. arXiv:1210.7449, doi:10.1016/j.physletb.2013.05.068.

[22] A. Andronic, P. Braun-Munzinger, K. Redlich, J. Stachel, Statistical hadronization of heavy quarks in ultra-relativistic nucleus-nucleus collisions, Nucl. Phys. A789 (2007) 334–356. arXiv:nucl-th/0611023, doi:10.1016/j.nuclphysa.2007.02.013.

[23] A. Kostyuk, Double, triple and hidden charm production in the statistical coalescence modelarXiv:nucl-th/0502005.

[24] Y. Liu, C. Greiner, A. Kostyuk, B_c meson enhancement and the momentum dependence in Pb+Pb collisions at LHC energy, Phys. Rev. C87 (2013) 014910. arXiv:1207.2366, doi:10.1103/PhysRevC.87.014910.

[25] L. Grandchamp, R. Rapp, Charmonium suppression and regeneration from SPS to RHIC, Nucl. Phys. A709 (2002) 415–439. arXiv:hep-ph/0205305, doi:10.1016/S0375-9474(02)01027-8.

[26] X. Zhao, R. Rapp, Transverse Momentum Spectra of J/ψ in Heavy-Ion Collisions, Phys. Lett. B664 (2008) 253–257. arXiv:0712.2407, doi:10.1016/j.physletb.2008.03.068.

[27] T. Song, K. C. Han, C. M. Ko, Charmonium production from nonequilibrium charm and anticharm quarks in quark-gluon plasma, Phys. Rev. C85 (2012) 054905. arXiv:1203.2964, doi:10.1103/PhysRevC.85.054905.

[28] Y. Liu, C. M. Ko, F. Li, Heavy quark correlations and the effective volume for quarkonia productionarXiv:1406.6648.

[29] B. Abelev, et al., Inclusive J/ψ production in pp collisions at $\sqrt{s}$ = 2.76 TeV, Phys. Lett. B718 (2012) 295–306. arXiv:1203.3641, doi:10.1016/j.physletb.2012.10.078.

[30] K. Aamodt, et al., Rapidity and transverse momentum dependence of inclusive J/psi production in pp collisions at $\sqrt{s}$ = 7 TeV, Phys. Lett. B704 (2011) 442–455. arXiv:1105.0380, doi:10.1016/j.physletb.2011.09.054,10.1016/j.physletb.2012.10.0

[31] F. Bossu, Z. C. del Valle, A. de Falco, M. Gagliardi, S. Grigoryan, et al., Phenomenological interpolation of the inclusive J/psi cross section to proton-proton collisions at 2.76 TeV and 5.5 TeVarXiv:1103.2394.

[32] Y. Liu, Z. Qu, N. Xu, P. Zhuang, Rapidity Dependence of J/psi Production at RHIC and LHC, J. Phys. G37 (2010) 075110. arXiv:0907.2723, doi:10.1088/0954-3899/37/7/075110.

[33] J. L. Albacete, et al., Predictions for p+Pb Collisions at sqrt s_NN = 5 TeV, Int. J. Mod. Phys. E Vol. 22 (2013) 1330007. arXiv:1301.3395.

[34] X.-N. Wang, F. Yuan, Azimuthal asymmetry of J / psi suppression in noncentral heavy ion collisions, Phys. Lett. B540 (2002) 62–67. arXiv:nucl-th/0202018, doi:10.1016/S0370-2693(02)02121-4.

[35] F. Arleo, P. Gossiaux, T. Gousset, J. Aichelin, Heavy quarkonium hadron cross-section in QCD at leading twist, Phys. Rev. D65 (2002) 014005. arXiv:hep-ph/0102095, doi:10.1103/PhysRevD.65.014005.

[36] A. Zoccoli, et al., Charm, beauty and charmonium production at HERA-B, Eur. Phys. J. C43 (2005) 179–186. doi:10.1140/epjc/s2005-02308-8.

[37] J. Beringer, et al., Review of Particle Physics (RPP), Phys. Rev. D86 (2012) 010001. doi:10.1103/PhysRevD.86.010001.

[38] B. Chen, Y. Liu, K. Zhou, P. Zhuang, ψ' Production and B Decay in Heavy Ion Collisions at LHC, Phys. Lett. B726 (2013) 725–728. arXiv:1306.5032, doi:10.1016/j.physletb.2013.09.036.